Fluid Phase Equilibria, 38 (1987) 307–308
Elsevier Science Publishers B.V., Amsterdam – Printed in The Netherlands

Contents Volume 38 (1987)

Volume 38, Nos. 1–2

Excess enthalpies of 1-iodoalkane + n-alkane mixtures. Measurement and analysis in terms of group contributions (DISQUAC)
J. Muñoz Embid, S. Otín, I. Velasco, C. Gutiérrez Losa (Zaragoza, Spain)
H.V. Kehiaian (Paris, France) 1–17

Equation of state for solutions of electrolytes in mixed solvents
W. Raatschen, A.H. Harvey and J.M. Prausnitz (Berkeley, CA, U.S.A.) 19–38

Local composition models for fluid mixtures over a wide density range
K.P. Johnston, S. Kim and J.M. Wong (Austin, TX, U.S.A.) 39–62

Vapor–liquid equilibria of strong electrolyte aqueous solutions
T. Sako, T. Hakuta and H. Yoshitome (Ibaraki, Japan) 63–81

The measurement and interpretation of the fluid-phase equilibria of a normal fluid in a hydrogen bonding solvent: the methane–methanol system
J.H. Hong, P.V. Malone, M.D. Jett and R. Kobayashi (Houston, TX, U.S.A.) 83–96

Vapour pressure data for 1-butanol, cumene, n-octane and n-decane and their statistically consistent reduction with the Antoine equation
J. Gregorowicz, K. Kiciak and S. Malanowski (Warsaw, Poland) 97–107

High pressure liquid–liquid equilibria for the system of water, ethanol and 1,1-difluoroethane at 323.2 K
T. Nakayama, H. Sagara (Yokohama, Japan) and K. Arai, S. Saito (Sendai, Japan) 109–127

Vapor pressures and osmotic coefficients: electrolyte solutions of methanol
P. Tomasula, G.J. Czerwienski and D. Tassios (Newark, NJ, U.S.A.) 129–153

Short Communication

Isobaric vapour–liquid equilibria for the ternary mixture: methyl ethyl ketone–water–toluene
B.S. Rawat, S.K. Gupta, S.M. Nanoti and R. Krishna (Dehra Dun, India) ... 155–161

Review

Heats of mixing in the critical region
J.J. Christensen, R.M. Izatt (Provo, UT, U.S.A.) and D.M. Zebolsky (Omaha, NE, U.S.A.) 163–193

Letter to the Editor

A cost-benefit analysis of the industrial use of a phase equilibria prediction model based on the perturbation theory
K.E. Gubbins and P. Clancy (Ithaca, NY, U.S.A.) 195–197

Erratum 199

Volume 38, No. 3

Thermodynamic properties of polar fluids from a perturbed-dipolar-hard-sphere equation of state
P.F. Bryan and J.M. Prausnitz (Berkeley, CA, U.S.A.) 201–216

Representation of the vapor–liquid equilibrium of the ternary system carbon dioxide–propane–methanol and its binaries with a cubic equation of state: a new mixing rule
J. Schwartzentruber, F. Galivel-Solastiouk and H. Renon (Paris, France) 217–226

Thermodynamics of mixtures containing bromoalkanes. I. Estimation of DISQUAC interchange energy parameters for 1-bromoalkane + n-alkane mixtures. Comparison with other haloalkanes
H.V. Kehiaian and I. Velasco (Paris, France) 227–244

Phase behavior of aqueous two-polymer systems
C.H. Kang and S.I. Sandler (Newark, DE, U.S.A.) 245–272

Excess molar volumes of carboxylic acid mixtures: binary mixtures of formic, acetic, propionic, n-pentanoic and iso-pentanoic acids at 298.15 K
A. Apelblat and E. Manzurola (Beer Sheva, Israel) 273–290

Volumes of mixing of decane isomers with normal hexadecane at 298.15 K. An interpretation in terms of the Van–Patterson theory
A.M. Awwad, F.A. Hassan and M.A. Salman (Baghdad, Iraq) 291–298

Short Communication

Excess volumes of mixing of binary systems of methanol with acetone and chloroform at 298.15 K.
S.E.M. Hamam and T.A. Al-Sahhaf (Safat, Kuwait) 299–303

Author Index 305

Contents of Volume 38 (1987) 307